	T	C D 1 -	
	Emergency		
	American Christia	an Academy	
	FOOD ALI	LERGY	
	School Year:	/	
Student:	Grade: Home	eroom Teacher:	DOB:
Asthmatic: 🗖 Yes 📮 No (i	increased risk for severe reaction)		
Allergen(s):			
Mother:	Home #:	Work #:	Cell #:
Father:	Home #:	Work #:	Cell #:
Emergency Contact:	Relatio	onship:	Phone:
	isea, abdominal cramps, vomiting, dia rtness of breath, repetitive cough, wh		
 LUNG Sho HEART "The severity of symptreatment is given i TREATMENT: Rinse control Then, call: Mother Father ACTION FOR MINOR RES 	rtness of breath, repetitive cough, wh ready pulse", "passing out" ptoms can change quickly – it : mmediately. ntact area with water if appropriate. Emergency Contact EACTION	arrhea neezing is important that Healthcare Provider	
 LUNG Sho HEART "The severity of symptreatment is given i TREATMENT: Rinse control Then, call: Mother Father ACTION FOR MINOR RES 	rtness of breath, repetitive cough, wh ready pulse", "passing out" aptoms can change quickly – it : mmediately. ntact area with water if appropriate.	arrhea neezing is important that Healthcare Provider	
 LUNG Sho HEART "The severity of symtreatment is given i TREATMENT: Rinse control of the severity of symtreatment severity of symtreatment severity of symtreatment severity of severity of severity of symtreatment severity of severit	rtness of breath, repetitive cough, wh ready pulse", "passing out" ptoms can change quickly – it : mmediately. Intact area with water if appropriate. Emergency Contact EACTION I I with symptoms I without w re:	arrhea neezing is important that Healthcare Provider waiting for symptoms	(medication/dose/route).
 LUNG Sho HEART "The severity of symthetic treatment is given in the severity of symthetic treatment is given in the severity of symthetic treatment is given in the severity of symthetic treatment should be initiated to the symphones (s) a give	rtness of breath, repetitive cough, wh ready pulse", "passing out" ptoms can change quickly – it : mmediately. ntact area with water if appropriate. Emergency Contact EACTION With symptoms I without w	arrhea neezing is important that Healthcare Provider waiting for symptoms	(medication/dose/route).
 LUNG Sho HEART "The severity of symtheter treatment is given in treatment is given in treatment is given in treatment is given in the should be initiated. Mother Instant Father ACTION FOR MINOR RING (s) a give	rtness of breath, repetitive cough, wh ready pulse", "passing out" aptoms can change quickly – it : mmediately. Intact area with water if appropriate. Emergency Contact EACTION I I with symptoms I without w re: e within 10 minutes or worsens, follow	arrhea neezing is important that lealthcare Provider waiting for symptoms	(medication/dose/route).
 LUNG Sho HEART "The severity of symtreatment is given in treatment is given in treatment is given in treatment is given in the severity of symtreatment is given in the severity of severe severe	rtness of breath, repetitive cough, wheready pulse", "passing out" ptoms can change quickly – it is mmediately. ntact area with water if appropriate. EACTION EACTION e within 10 minutes or worsens, follo EACTION EACTION	arrhea neezing is important that I Healthcare Provider waiting for symptoms ow steps for Major Reaction	(medication/dose/route). 1 below. MPTOMS ARE PRESENT, DIATELY AND 911 WILL BE
 LUNG Sho HEART "The severity of symple treatment is given in treatment is given in treatment is given in treatment is given in the severity of symplement should be initiated in the only symplement should be initiated be initiated should be initiated	rtness of breath, repetitive cough, wheready pulse", "passing out" aptoms can change quickly – it a mmediately. Intact area with water if appropriate. EACTION I I With symptoms I without water if within 10 minutes or worsens, follo EACTION S I No Special instructions:	arrhea neezing is important that Healthcare Provider waiting for symptoms ow steps for Major Reaction LERGEN OCCURS, SY VILL BE GIVEN IMME PORTED TO NEARES' phrine, a student may feel d	(medication/dose/route). n below. MPTOMS ARE PRESENT, DIATELY AND 911 WILL BE T HOSPITAL. izzy or have an increased heart
 LUNG Sho HEART "The severity of symthese treatment is given in treatment is given in treatment is given in treatment is given in the should be initiated. Mother I Father ACTION FOR MINOR RI Treatment should be initiated. If the only symptoms (s) a give	rtness of breath, repetitive cough, wheready pulse", "passing out" ptoms can change quickly – it is mmediately. Intact area with water if appropriate. EACTION I INDUCTION I INDUCTION INDUCTION INDUCTIONIAL INDUCTIONIALIAL INDUCTIONIALIAL INDUCTIONIALIALIALIAL INDUCTIONIALIALIALIALIALIALIALIALIALIALIA	arrhea neezing is important that Healthcare Provider waiting for symptoms ow steps for Major Reaction ERGEN OCCURS, SY VILL BE GIVEN IMMET PORTED TO NEARES' phrine, a student may feel d hould be transported to the	(medication/dose/route). n below. MPTOMS ARE PRESENT, DIATELY AND 911 WILL BE T HOSPITAL. izzy or have an increased heart hospital by ambulance

This plan is in effect for the current school year and summer school as needed.